

THE BIOGEOCHEMISTRY OF
THE AMAZON BASIN

Index

- abandoned pasture, 143
 ABLE experiments, 29, 50
 aboveground biomass, 7, 229
 carbon loss due to deforestation, 86
 nutrient stocks, 145
 ABRACOS, 36
 acetate, 49
 Acre, 125
 Acri-Ferrasol group, 55
 adaptive strategies, 11
 aerosols, 5, 44
 natural biogenic component, 46
 long distance travel, 47
 aerosol mass, 44
 aerosol particles,
 vegetation as source, 44
Aeschynomene sensitiva, 215
 Africa, 43, 48, 68
 Africanization, 122
 ageotropism, 61
 Agroextractive Cooperative of
 Xapuri(CAEX), 133
 agroforestry, 106, 114, 117
 agrosilvopastoral systems, 106
 air temperature, 34
 positive trend, 37
 albedo, 35
 Alfisols, 87, 165
 algae, 45, 211, 235
 algal chlorophyll, 263
 Ali-acrisols, 55, 62
 Ali-ferrasols, 62
 alkalinity, 286
 as a hydrological tracer, 253
 allochthonous carbon, 253
 alluminum-accumulating species, 78
 Alta Floresta, 46
 Altamira, 34
Alternanthera pilosa, 248
 Aluminum, 54, 201
 Amapá state, 108, 323
 AmasSeds Project, 329,336-337, 352
 Amazon,
 central, 10, 21
 eastern, 8, 21, 25
 western, 21, 25
 Amazonian hydrologic cycle, 29
 Amazon shelf, 14
 carbon cycling, 345
 nutrient trap behavior, 332
 oxygen data, 335
 processes controlling biogeo-
 chemical reactions, 336
 sedimentary record, 329
 sources of POC, 334
 Amazon trough, 308
 amerindian activities, 122
 amino acids, 13, 294, 297
 ammonium uptake, 64
 anaerobic environments, 200
 anatase, 309
 Anavilhanas archipelago, 259
 Andean Amazon, 5
 Andean Cordillera, 188, 314
 Andean glaciers, 187
 Andes, 13, 53, 87, 225, 278, 290, 309
 upper levels, 18
Andropogon gayanus, 89
 anion exchange capacity, 63
 annual duration of inundation, 211
 ant nests,
 enriched in nutrients, 152
 ants,
 reduce the resistance to penetration of
 deep, intact soil, 152
 seedling-eating, 142
 Cutter, 152
 aquatic macrophytes, 211
 Araraquara, 169
 Argentina,
 northern, 37
 Ariquemes, Rondonia, 180
 Arsenic, 14, 307, 312
 assimilation capacity, 10
 Atlantic forest region of Bahia, 112
 Atlantic ocean, 4, 188, 275, 325
 western, 14
 atmospheric deposition, 259
 atmospheric transport, 5
 Atta sexdens, 152
 Aquatic Terrestrial Transition Zone
 (ATTZ), 210, 217, 219, 236, 237, 261
 austral warm season, 18
 Australia, 68
 authigenic minerals,
 in Amazon sediments, 353
 Azougue (liquid mercury), 265

 Bacia Modelo, 194
 bacteria, 44, 197
 imperfecti, 197
 Bahia, 118
 B-alanine, 297
 barium, 312
 Barro Branco stream, 10
 Barro Colorado, 64
 base cations, 55
 sum of, remain elevated in pasture soils, 97

Batrachospermum, 198
 BATS, 35
 Bauxite mining, 113
 Bauxite tailings, 264
 Belém, 141, 225
 air temperature, 34
 Belém Brasília highway, 88
 benthic diatoms, 198
Bertholletia excelsa, 107, 131
 biochemical oxygen demand, 202
 biodegradation, 275
 biogenic silica, 329
 burial of, 332
 biogeochemical fluxes,
 regionalization of, 267
 biological production on shelf, 14
 biomass, 11
 aboveground, 56, 57, 86, 172
 belowground, 56, 57
 burning, 42, 43, 49, 86
 cerrado, 73, 75
 cerrado compared to terra firme forest, 74
 herbaceous layer cerrado, 76
 nonforest vegetation, 178
 post-logging, 174
 Savanna, 179
 biomass accumulation, 139
 rate of, 140, 142
 biomass development, 54
 biomass estimates,
 direct methods, 171
 how to improve, 179
 indirect methods, 171
 should prioritize the measure of big
 trees, 181
Bixa orellana, 112
 black carbon, 46
 black water rivers, 11, 188, 278
 Bolivia, 323
 Altiplano, 18
 Bolivian high, 18
Brachiaria brizantha, 86, 90-92, 95, 97-99,
 122, 146
Brachiaria decumbens, 91, 99
Brachiaria humidicola, 92, 99
 Bragantine Zone, 141
 Branco river, 199, 314
 Brazil,
 state of Roraima, 28
 northeast, 37
 southern, 37
 Brazilian Institute of Geography and
 Statistics (IBGE), 177
 Brazilian shield, 308

- Brazil nut, 131
 buffering capacity of Amazon stream systems, 10
 Buritizaís, 69
- C-4 grasses, 215
 C-14, 334
 changes in, 299
 Cadmium, 202, 307
Cajanus Cajan, 162
 Calanoid, 255
 Calcifugous or Calcicole species, 72
 calcium, 53-54, 79, 97, 189, 194, 201
 foliar concentration, 226
 immobilized in slash, 98
 in herbaceous plants, 216
 limitation, 6, 62
 losses from the system due to combustion of biomass, 86
 retained on the soil exchange sites, 145
 stocks, 109
Calopogonium spp., 93
 Campo limpo, 69
 Campo sujo, 72
 CAMREX Project, 277, 287, 335
 Caquetá, 55, 61
 Carajás area, 198
 carbohydrates, 13, 294
 carbon, 236
 allocation patterns, 119
 cycles of, 3, 7, 42
 increase soil C, 112
 loss during conversion, 145
 losses from the system due to combustion of biomass, 86
 quantity stored in soils and biota, 9, 165
 carbon dioxide, 42, 165
 a sink for atmospheric, 4, 9, 43
 concentration, 287
 emission rates, 257
 exchange with atmosphere, 237
 production, 280
 released to the atmosphere, 8
 temperature increase, 36
 source of, 43
 carbon dynamics,
 conceptual model in a forested landscape, 126
 floodplain lakes, 246
 carbon flux measurements, 42
 carbon in soil,
 for the Legal Amazon Basin, 166
 carbon sequestration, 142
 carbon stocks, 165
 cation exchange capacity, 63, 97, 225
 cattle, 7
 cattle grazing,
 affects the rate of P movement and distribution, 90
Cecropia palmata, 99
Cecropia spp., 223
Cedrela odorata, 108, 111
Ceiba pentandra, 222, 225
 Cerradão, 72
 Cerrado, 7, 43, 68, 72
 biomass, 73, 75
 geographic distribution of species, 73
 nutrient stocks, 74
Centrosema spp., 93
 Characiform detritivores, 255
 Chico Mendes, 129
 Chico Mendes Extractive Reserve, 125, 129, 134
 chlorine, 201
 chronosequence technique, 140
 Cinnamyl phenols, 296
 Cladocera, 255
 clay minerals, 54
 cleared forest,
 deeper convective boundary, 36
 clear water rivers, 278
 climate change, 9, 11
 climate system,
 global, 5, 17
 climate variability,
 low-frequency variation, 31
 cloud condensation nuclei, 46
 over Amazonia and Africa, 47
 cloud cover, 36
 cloud formation mechanisms, 44
 coarse particulate C and N,
 relationships between and river slope, 290
 coastal waters,
 oxygen distributions, 328
 cobalt, 307
 COLA GCM, 35
 Colombia, 60, 61, 169, 309, 323
 colonists, 6
 Congo, 275
 continental shelf, 328
 convection,
 land-based, 17
 copper, 12, 202, 307, 312
Cordia multispicata, 99, 112, 143
 Costa Rica, 50, 64, 95, 112, 200
 Cretaceous, 69, 70
 chromium, 202
 crops, 11
 sustainability, 65
 cross-equatorial flow,
 northern, 20
 Cuiabá, 46
 cyclopoid copepods, 255
- dams, 276
 Daniel Ludwig, 108, 156
Davilla kunthii, 150
 decomposition bags, 62
 deep roots, 8
 deep root systems,
 re-establishment of, 150
 deforestation, 7, 9, 12, 17, 84
 causes increase in erosion, 202
 dependance on albedo, 35
 essential element in reducing, 6
 impact of in situ, 35
 reduction by extractive reserves, 127
 regional climate, 35
 denitrification, 93, 258
 in bacteria, 95
 deposition/resuspension cycle, 323
Desmodium spp., 93
 detrital food chain, 197
 development pressures, 14
- diatom community, 14
 direct litterfall, 13
 direct nutrient cycling, 144, 189
 discharge of industrial and human waste, 202
 dish-shaped lakes, 239
 dissolved humic compounds, 199
 dissolved organic carbon, 280
 riverine, 254
 dissolved organic compounds, 49, 193, 201
 dissolved organic nitrogen,
 flux of riverine, 342
 dystrophic soils, 68
 droughts,
 midsummer, 24,
 fires, 27
Drulia braunii, 198
 Ducke Reserve, 189, 192, 195
- Echinochloa polystachya*, 11, 212, 214-215, 249, 255, 295
 ectorganic horizon, 61
 ECMWF, 29
 Ecuador, 323
Eichhornia crassipes, 212, 214, 264
 elemental ratios,
 carbon, 13
 El Niño Southern Oscillation (ENSO), 5, 18, 25, 31, 122, 242
 EMBRAPA, 111, 118, 133
 emissions,
 natural biogenic, 47
 pyrogenic, 47
 Entisols, 59, 69, 165
 EOS-DNAEE network, 28
Eperua rubiginosa, 55
 erosional increase,
 due to deforestation, 202
Eucalyptus spp., 108
Eucalyptus urogandis, 110
Eugenia inundata, 211
Eunotia, 198
 evapoconcentration effect, 148
 evaporation,
 local, 17
 evapotranspiration, 201
 annual ratio, 28
 estimates, 29
 potential, 71
 exchangeable nutrient patterns, 56
 extractive reserves, 8, 122
 appropriate conditions, 125
 FAO, 165, 173
 fauna, 4
 Fazenda Agua Limpa, 73
 Fazenda Vitoria, 87, 94, 142
 feces, 7
 Ferralsols, 54-55
 fertilization, 7, 65
 response to, 143
 fertilizers, 9
 FGGE, 29
 fine roots,
 surface mats or, 6
 fine-root biomass,
 distribution of in the secondary forest, 147
 fire, 7, 27, 37

- fuel material, 79
- adaptive strategies, 79
- fish, 9, 11, 276
 - carbon sources for adult, 255
 - carbon source for larval, 255
 - herbivorous and omnivorous, 256
- fisheries, 4
- floating meadow community, 236
- flood cycles, 211
- flooded forest,
 - contribution to floodplain net production, 250
- flooding, 12
- floodplain, 10
 - herbaceous communities, 11
- floodplain forests, 11, 252
 - floristic composition of, 219
 - net production estimates for, 250
- floodplain lakes, 11, 236
 - carbon dynamics, 246
 - cation fluxes in, 236
 - flow paths of inflows to, 243
 - human disturbances, 264
 - impacts of deforestation on, 266
 - light attenuation in, 241
 - net production rates, 252
 - number along mainstem, 236
 - primary production in, 246, 246
 - vertical mixing, 244
 - water balance of, 242
- floodplain systems, 209
 - productivity of, 210
- flood pulse, 209, 224
- flood stress, 224
- floodwaters, 11
- flood waves,
 - passage of, 235
- flora, 4
 - adaptive mechanisms, 3
- floristic composition,
 - Cerradões and cerrados, 72
- FLUAMAZON, 29
- Fly river, 351
- foodweb, 12
- Fordlandia, 107
- forest adaptation, 64
- forest biomass,
 - estimation methods for, 170
- forest burning,
 - sources of metals, 324
- forest clearing,
 - loss of aboveground nutrients during, 144
- forest flammability, 150
- forest-to-pasture conversion, 84, 86
- forests,
 - aerosol emission, 42
 - area converted to pasture, 84
 - burning, 125
 - nonflooded, 6
 - regeneration, 8
 - rate of recovery of secondary, 8
 - fires, 27, 37
 - terra firme, 53
 - Diptercarp, 56
 - Caatinga, 57
 - Tall Amazon Caatinga, 57
 - trace gas emission, 42
 - rate of recovery, 58
- Formate, 50
- FPOC,
 - Andean origin, 291
- fragens, 20
- fucose, 296
- fulvic acids, 199
- fungal hyphae, 147
- fungi, 45
 - gallery forests, 68-70
 - G-aminobutyric acid, 297
 - Ganges, 351
 - Garimpos, 323
 - gas exchange, 275, 291
 - general circulation models(GCM), 34
 - atmospheric, 35
 - grid resolutions, 34
 - generation of hydroelectricity, 37
 - geomorphology, 11
 - Global Positioning System(GPS), 134
 - Gmelina arborea*, 108, 156
 - gold, 198
 - Gran Chaco, 18
 - Grande Carajás Program, 110
 - greenhouse gases, 4
 - groundwater, 10
 - Guadua* sp., 129
 - Guttation, 45
 - Guyana, 55, 60
 - Guyana shield, 308
- heavy metals, 10
- Henry Ford, 107
- herbaceous C4 plants, 230
- herbaceous macrophyte, 238
 - contribution to floodplain net production, 249
- herbaceous plants,
 - indicators of ecological conditions, 212
- number of species, 213
- herbaceous vegetation,
 - biomass production of, 214
- heterotrophic,
 - metabolism, 12
 - activity, 14
- Hevea brasiliensis*, 107
- HiBAM project, 308
- Hoplosternum littorale*, 202
- Huanghe river, 351
- human activity, 29-30
- Humboldt and Bonpland, 132
- humic acids,
 - high molecular weight, 334
- humid forest, 30
- humidity deficit,
 - pasture area, 98
- hydrocarbons,
 - photochemical oxidation of, 44
- hydro-electric dams, 12, 176
 - impacts of, 264
- hydrological cycle, 210
- hydrological functions,
 - recovery of, 150
- hydrological modeling, 34
- hydrological recovery, 139
- hydrolyzable amino acids, 298
- hydrometeorological conditions, 26
- hydrophobic sorption, 299
- Hymenachne amplexicaulis*, 215
- Hyparrhenia rufa*, 92
- Hyphmycete fungi, 197
- Hypsipyla grandella*, 108, 111
- IBAMA, 125, 133, 174
- Icá river, 188, 309
- Igapo, 11, 212
 - diversity in, 224
 - Igapó trees,
 - nutrient use efficiency of, 226
- Ilha de Marchantaria, 219
- Inceptisols, 69, 165
- INCRA, 125
- industrial expansion, 42
- Inga affinis*, 112
- Instituto Nacional de Pesquisas da Amazônia(INPA), 188
- Instituto Nacional de Pesquisas Espaciais, (INPE), 34
- inundation,
 - annual duration of, 211
- invertebrate scrapers, 197
- ion-depleted waters, 200
- Iquitos, 27, 111
- Iron, 54, 201
- Iron oxides, 314
- irrigation, 4
- Isoprenes, 42
- isotope techniques, 170
- isotopic tracers, 147
- ITCZ, 31
- Japura River, 188, 309-310
- Jarí Forestal e Agropecuaria, 107
- Jarí, 180
- Jarí Project, 110, 156
- Jarí River, 108, 156
- Ji-Paraná, 98
- Kaolinsols, 54
- kaolinite, 55
- kaolinite-organic matter complexes, 199
- kaolinitic soils, 97
- King Ranch of Texas, 157
- La Cuenca, 192, 194
- Lago Calado, 192, 194, 195
- lake,
 - Batata, 264
 - Calado, 235, 242, 247, 258
 - Camaleão, 217, 235,236, 251, 260
 - Cristalino, 266, 310
 - Janauaca, 254
- lake morphology, 238
- land abandonment, 8
- land clearing, 85
- land conversion,
 - net effects on nutrient inputs to streams, 201
- LANDSAT images, 34, 172, 243-244
- land surface,
 - lost to the system, 28
- land use, intensification of, 158
- La Niña, 18, 25

- Large Scale Biosphere-Atmosphere (LBA), experiment in Amazonia, 37
- large-scale plantations, economic sustainability of, 108
- La Selva Biological Station, 110, 112
- lateritic nodules, 58
- lateral-levee lakes, 239
- lateral subsurface flowpaths, 191
- Latossolo Amarelo, 165
- Latossolo Vermelho Amarelo, 165
- Latossolo Vermelho Escuro, 72
- Latosols, 69, 165
- leaching processes, 6
- lead, 307
- released as dissolved form, 324
- Lecythis idatimon*, 150
- Leersia hexandra*, 248
- leguminous species, 72, 93
- lignin, 294, 299
- lignin molecules, 13
- limestone outcrops, 70
- litter blow-in, 186
- litterfall, 58
- decomposition of, 260
- rates, 190
- litter production, cerrado, 77
- Llanos region, 193
- macrophytes, 11
- Madeira river, 188, 319
- magnesium, 53-54, 97, 189, 194, 201
- limited, 6, 62
- deficiency, 62
- foliar concentration, 226
- in herbaceous plants, 216
- retained on the soil exchange sites, 145
- stocks, 109
- mainstem rivers, 12, 13
- major tributaries, 13
- Malaysia, 129
- Manacapuru, 34
- Manaus, 24, 32, 50, 99, 180, 189, 191, 202, 213, 235, 237, 241, 277, 308
- harbor, 209
- manganese, 14, 201, 307
- Marabá, 100
- Maracá, 59, 60
- Maranõn, 309
- Marchantaria, 226, 277
- marine influences, 50
- marine nitrogen, particulate organic, 343
- Mato Grosso state, 46
- Mauritia vinifera*, 69
- mercury, 203, 276, 307
- liquid, 265
- methylation and bioaccumulation of, 266
- pollution, 12
- mercury contamination, 265
- Mesophytic forests, 69
- metamorphism, 70
- methane, 4, 11, 42, 165, 186, 235, 256, 276
- emission rates, 43
- exchanges with atmosphere, 237
- fluxes, 200
- methyl bromide, 44
- methyl chloride, 44
- microclimatic conditions, 100
- microbial oxidation, 328
- mining, 12, 276
- activities, 80
- Mississippi delta, 351
- mahogany, 118
- moisture convection, 35
- moisture convergence, percent of precipitation, 29
- moisture sink, 30
- molybdenum, 307
- monospecific plantations, 113
- Monte Dourado Forestry Company, 108
- mulch decomposition rate, 113
- mycorrhizae fungi, 6, 9, 113
- mycorrhizal association, 78, 147
- mycorrhizal symbiosis, 63
- Myrcia sp.*, 211
- myth of the Amazon as a source of unbri-dled productivity, 157
- NAR, 31
- NASA, 324
- NASA-DAO, 29
- National Council of Rubber Tappers, 134
- National Institute for Amazonian Research, 110
- Navicula*, 198
- NCEP, 29
- Negro river, 13, 188, 198, 259, 309
- neophytes, introduction of, 213
- net primary productivity (NPP), 216, 230
- nickel, 14, 307
- nitrate, 201, 328
- distributions, 291
- first-order streams draining pastures, 96
- nitricification, 64, 93, 95, 258
- in pastures, 7
- net, 96
- nitrogen, cycles of, 3, 7
- dissolved organic flux to shelf, 342
- foliar concentration, 226
- increased soil N, 112
- in herbaceous plants, 216
- loss during conversion, 145
- losses from the system due to combus-tion of biomass, 86
- lower availability in old pastures, 96
- mobilized annually from deforestation, 93
- moderate acidity, 55
- net uptake, 218
- pools, 11
- recovery, 65
- returned in urine, 95
- total flux to shelf, 343
- use efficiency, 60
- nitrogen budget, water column recycling, 344
- nitrogen deficiency, 94, 262
- nitrogen fixation, 77, 78, 94, 258, 264, 352
- contribution by mycorrhizae, 227
- in the grass rhizosphere, 93
- rates on the Amazon shelf, 344
- nitrogen mineralization in pastures, 7
- rates, 63
- nitrogen transformations, 266
- nitrogen turnover, 261
- nitrogenous organic substances, sorption on clay minerals, 298
- nitrous oxide, 4, 7, 42, 165
- flux, 94
- increases in emissions, 95
- N₂, production rates, 345
- N₂O, 7
- flux, 94, 200
- increases in emissions, 95
- nonmethane hydrocarbons (NMHC), source of, 44
- nontimber forest product extraction, 123
- Normalized difference vegetation index (NDVI), 172
- nutrient, accumulation, 139
- conserving mechanisms, 9
- efficient recycling of, 156
- in storm water, 190
- leaching of, 53
- recycling, 54
- recycling mechanisms, 162
- sequestration of, 217
- stocks in cerrado, 74
- nutrient availability, 54, 56
- nutrient cycling, indicators, 65
- nutrient fluxes, modeling of, 217
- nutrient limitation, 263
- nutrient losses, due to removal of wood, 110
- effects of perturbation on, 107
- nutrient pools, combustible components, 79
- nutrient stocks, 144
- nutrient stress, 71
- nutrient trap, behavior on the Amazon shelf, 332
- Nutrient Use Efficiency (NUE), 60, 114, 116
- Óbidos, 25, 319, 277
- organic carbon, buried in shelf sediments, 347
- coarse particulate, 281
- deposits on the floodplain, 258
- fine particulate, 281
- total balance for lakes, 258
- organic matter, budget, 58
- degradation history, 296
- dynamics and pools, 13
- management of, 65
- marine burial of, 328
- mineralization, 58
- oxidation, 352
- partitioning between fine particulate and dissolved forms, 293
- sources, 294
- organic molecules, nitrogen-rich, 302
- organic residues, 64

- organic wastes, 10
 Orinoco basin, 53
 Oryza, 212
 overgrazing, 88
 overland flow, 10
 oxbow lakes, 239
 oxidation/reduction, 275
 Oxisols, 54, 59, 69, 85, 87, 89, 156, 165, 170, 192, 194, 226
 oxygen,
 cycles of, 3
 production on the shelf, 349
 riverine flux to shelf, 347
 oxygen consumption,
 for the shelf, 347
 oxygen deficiency, 211
 ozone,
 production or destruction, 44
 concentrations, 44
 ozone budget, 44
 Pacific Ocean, 49
 Paleozoic, 308
 Pampean Sierras, 18
 Panama, 64
Panicum Maximum, 85, 99
 Pantanal,
 GCM grid resolutions, 34
 Pará, Brazil, 96, 140, 156
Paragominas, 87, 94, 111, 141, 148, 158, 180
Parapiptadenia pterosperma, 112
 participatory research, 131, 135
 particulate phosphorous,
 transfer in the estuary, 332
Paspalum fasciculatum, 214–216
Paspalum repens, 214, 295
 pasture degradation, 84
 by grazing, 158
 pastures, 6, 11
 abandonment, 7, 143
 fertility, 7
 invaded by dicotyledonous species, 99
 overgrazing, 98
 productivity, 7, 84
 pasture soils,
 retention of base cations, 98
 Patch Dynamics Concept, 188
 Patos de Minas, 70
 periphyton, 11, 211
 C-13 of, 255
 contribution to floodplain net
 production, 251
 Peru, 169, 309, 323
 Peruvian-Bolivian Altiplano, 18
 PESACRE (Agroforestry Systems Research
 and Extension Group of Acre), 133
 pest and disease problem, 119
 pesticides, 9
 P-32, 6, 63
 additions, 263
 pH, 192
 elevation in pasture soils, 97
 phosphate, 328
 uptake by fine roots, 6
 available, 55
 phosphorus, 54, 79, 236, 286
 advection of, 341
 availability, 84
 cycles of, 3
 deficiency, 57
 deposition rate, 341
 diffusive flux of, 342
 efficiency, 60
 fixing capabilities, 55
 foliar concentration, 226
 immobilization of, 92
 in aerosols, 46
 in herbaceous plants, 216
 in the mineral soil of the várzea, 225
 limitation, 263
 limited, 6, 62
 limiting factor to primary productivity, 88
 loss during conversion, 145
 losses from the system due to
 combustion of biomass, 86
 net flux to the ocean, 342
 net uptake, 218
 nutrition, 101
 P fluxes, 60
 particulate, 280
 potential soil sources, 145
 recovery, 65
 released to the mixing zone, 341
 soil chemistry, 225
 soil uptake, 92
 stocks of, 89, 109
 transferred in feces, 91
 transformation, 64
 phosphorous fixation, 63, 92
 organic acids prevent, 156
 photochemical oxidation of hydrocar-
 bons, 44
 photosynthesis, 42, 275, 291
 phytoplankton, 261
 gross production by, 247
 phytoplankton production, 247, 314
Pinus caribaea, 108
Pinus caribaea var. hondurensis, 108
 pioneer forest community, 223
Piranhea trifoliata, 222
Pistia stratiotes, 214
 placer mining, 203
 plagioclase, 314
 plankton, 11
 open-water, 14
 planktonic respiration, 254
 plant growth,
 nutrient limitations to, 55
 plantation management, 017
 plantation tree species, 106
Plathymenia foliolosa, 112
 Pleistocene, 69
 Podzólicos, 165
 Podzólico Vermelho Amarelo, 165
 pollen, 44
 polysaccharide loss, 301
 Porto Trombetas, 113
 Porto Velho, 34
 potassium, 54, 79, 97, 192, 201
 foliar concentration, 226
 in herbaceous plants, 216
 leached from plant leaves, 50
 limited, 62
 long term supply, 146
 loss during conversion, 145
 losses from the system due to combus-
 tion of biomass, 86
 net uptake, 218
 potential soil sources, 145
 recovers, 65
 potassium amendments, 97
 Precambrian, 70
 Precambrian Shield, 55, 188
 precipitation,
 local recycling, 17
 average, 28
 preservationist/developmentalist dichoto-
 my, 123
 primary production, 11, 345
 herbaceous layer of a cerrado, 75
Prochilodus nigricans, 255
 productivity,
 of Amazon forests, 6
 tropical grasses, 93
 protozoa, 45
Pseudobombax munguba, 260
 Purus river, 188
 pyrogenic emissions, 47

Qualea grandiflora, 78
Qualea multiflora, 78
Qualea parviflora, 78
 quartz, 54, 309
 Quartzipsamments, 70, 71
 Quaternary, 69

 RADAMBRASIL, 72, 166, 173, 174, 178, 237
 radiation budget, 42
 radiocarbon, 87
 rainfall,
 inter-annual variability, 20
 low, 18
 mean annual, 241
 mechanisms which produce, 21
 negative anomalies, 18, 25
 positive anomalies, 25
 variability in, 5, 17
 rare earth elements, 320
 rates of deforestation,
 decrease in precipitation due, 5
 increase in the Amazon basin, 30, 35
 decrease in the Amazon basin, 35
 increase surface temperature, 36
 decrease evapotranspiration, 36
 Recife, 63
 recycling,
 nutrients, 54
 precipitation and evaporation, 17
 rainfall in region, 30
 water in hydrological cycle, 28
 Redfield ratio, 337
 refractory material, 12
 regional chromatography, 13
 model, 302
 replacement of forest,
 by grassland, 36
 Reserva Ducke, 192, 195
 residual oil combustion, 324
 respiration, 42, 291
 rhamnose, 296

- rhizosphere, 63
 ria lakes, 239, 242
 riparian
 forests, 10
 plants, 198
 wetlands, 19
 riparian vegetation,
 aggrading, 202
 riparian zones,
 buffering, 202
 River Continuum Concept, 188
 river discharge, 28
 river-lake interface, 236
 river/ocean mixing, 328
 carbon transported into the Amazon, 334
 rivers,
 Guamá, 219
 Jari, 108
 Juruá, 219
 Madeira, 26, 34
 Napo, 219, 225
 Negro, 13, 22, 23, 26, 32, 34, 53
 Obidos, 25, 32
 Purus, 26
 Santarem, 25
 Solimões, 25, 34, 222
 Tapajós, 25, 118
 Ucayali, 111
 Xingu, 34, 219
 road construction, 80
 Rondonia, 93, 95, 96, 98, 168-169
 Ariquemes, 180
 pattern of vegetation, 34
 root biomass, 62
 root mat, 6, 61
 root systems,
 deep, 143
 Roraima, 61
 rubber extraction, 122
 rubber plantations, 129
 runoff,
 organic carbon in upland, 254
 surface, 185
 subsurface storm, 185

 Sahara dust, 48
Salix martiana, 211-212
Salvertia convallariodora, 78
Salvinia auriculata, 214
 San Carlos de Río Negro, 53, 57, 61, 62,
 64, 65, 201
Salix humboldtiana, 228
 Santarem/Cuiabá road, 118, 307
 São Paulo, 71, 77, 157
 SAR, 31
 savanna, 6, 222
 soils, 92
 savannization, 122
 SCAR-B experiment, 46
 seafloor, 14
 seasonal water shortages, 6
 seasons,
 austral warm, 18
 sea surface temperature (SST),
 tropical Atlantic, 17
 variations, 31
 dipole, 31

 secondary forest,
 areal coverage, 139
 distinctive phases, 140
 establishment of, 140
 growth rates, 65
 land use effects on mechanisms of
 establishment, 142
 sediment,
 exchanges between the floodplain and
 channel, 257
 sedimentary record,
 on the Amazon shelf, 329
 selective logging, 201
Semaprochilodus insignis, 255
 Serra do Navio, 323
 sewage inputs, 351
 shelf phytoplankton,
 sources of nutrients, 331
 shifting cultivation, 125, 156
 Silica, 14, 192
 biogenic, 329
 burial rate, 340
 offshore advective flux, 339
 silicate, 328
 diagenetic recycling, 339
 silviculture,
 plantations, 8
 SLAR, 237
 slash-burn, 64, 85, 140, 203
 agriculture, 84
 slash-burn areas,
 estimate the biomass volume within, 172
 Smectite-vermiculite, 314
 sodium, 194, 201
 in herbaceous plants, 216
 soil,
 deep, 144
 distribution in the cerrado region, 71
 fertility, 6, 54
 kaolinitic, 97
 sink or a weak source of CO₂, 43
 soil Carbon,
 problem of missing data, 169
 soil Carbon stocks,
 land use practices affect, 170
 soil erosion, 13
 rates of, 80
 soil fertility,
 gradients in, 70
 soil moisture,
 deep, 143
 soil organic matter, 56
 turnover of, 87
 residence time, 87
 soil organic Carbon, 165
 soil Phosphorous uptake, 91
 soil taxonomy, 71
 soil water,
 deep uptake, 150
 low availability of, 71
Solanum crinitum, 143
 Solimões River, 198, 309, 319
Sorghum arundinaceum, 248
 South Atlantic Convergence Zone (SACZ),
 20, 71
 Southern Oscillation (SO), 17
 signal of, 37

 species,
 richness composition, 55
 species composition, 11, 224
 spiraling effect, 197
 Spodosols, 59
 spores, 44
 stable isotope ratios,
 carbon, 13
Sterculia pruriens, 150
 stomatal conductance, 99
 stratospheric ozone depletion, 44
 streams, 10, 185
 strontium 86/87, 147, 312
 subsurface runoff, 13
 successional processes, 213
 sulfate, 50, 201, 310
 Superintendency for the Development of
 the Amazon (SUDAM), 157, 177
 suspended particulate material,
 downstream distributions, 281
 suspended sediment,
 estimated concentrations of, 244
 sustainability, 64
 sustainable development, 14, 15, 123
 sustainable plantations, 116
 sustainable systems of farming and
 forestry, 162
Swietenia macrophylla, 107, 111, 112, 118
Symmeria paniculata, 211
Synedra, 198

Tachigalia myrmecophila, 150
 Tapajós river, 118, 237, 309
 termites,
 role in decomposition, 77
 terpenes, 42
 terra firme, 156
 nonflooded,
 Tertiary, 69, 308
 thermodynamics, 29
 throughfall, 10, 50, 191
 timber plantations,
 small-scale, 119
 titanium, 198
 trace elements,
 temporal variability, 319
 transport of the particulate, 317
 trace metals, 13, 307
 abundances, 13
 anthropogenic and natural fluxes, 323
 pollution, 14
 transport of dissolved, 317
 transpiration, 45, 148
 transportation, 37
 tree litter, 113
 tree plantations, 107,
 contribute to soil restoration and faci-
 litate natural forest regeneration, 112
 tree-ring analysis, 228
 tree species,
 density, 55
 TRIMM-LBA, 21
 tritium-labeled bacteria, 254
 Trombetas, 180, 264, 309
 tropical Pacific, 36, 44
 tropical Atlantic, 36, 44
 tropical forests,

- conversion of - to cattle pastures, 93
- tropical plantations, 106
- tropospheric convergence,
 - upper and lower, 30
- tropospheric zonal temperature, 18
- Ucayali River, 111, 309
- Ultisols, 54, 59, 71, 85, 87, 156, 192, 194, 226
- upper continental crust, 317
- upper troposphere,
 - meridional flows, 18
- uranium, 312
- U.S. Soil Taxonomy, 165
- UV radiation, 43
- vanadium, 14, 312
- vanillic acid, 297
- vanillin, 297
- vanillyl phenols, 296
- variation between forest and agroecosystems,
 - carbon stocks, 126
- Vargem Grande, 277
- várzea, 11, 209, 222, 224, 275, 277, 300, 323
 - biomass of, 228
 - nitrogen fluxes in, 226
 - soil chemistry, 225
- várzea soils,
 - nutrient stocks (Table 13.8) in, 225
- vegetation,
 - floodpulse-induced zonation of, 211
- vegetation islands, 99
- Venezuela, 60
- veranico, 24, 36
- vesicular-arbuscular mycorrhizal fungi (VAM), 63
- Vochysia divergens*, 78
- Vochysia ferruginea*, 112
 - protect soil against erosion, 114
- Vochysia guatemalensis*,
 - Ca and Mg recycling, 114
- Vochysia thyrsoidea* Pohl., 78
- volatile organic compounds, 42
- water balance,
 - indirect methods, 28
- water column respiration,
 - in Amazon floodplain lakes, 252
- water management, 9
- water quality, 9, 12
 - threats, 4
- water table,
 - fluctuations of, 53, 72
- water vapor flux,
 - interannual variations, 29
- wet deposition, 5, 49
- wetland, 10
- wetland seepage, 186
- whitewater rivers, 11, 209, 278
- World Bank, 160
- Xapuri, 129, 132
- Xingú river, 237, 309
- Yurimaguas, 169
- Zaire, 351
- zebu cattle, 91
- zinc, 202, 307
- zircon, 198, 309